

Terrestrial & Palustrine Plant Communities Of Pennsylvania

By Jean Fike

Hemlock (white pine) - red oak - mixed hardwood forest

This type is similar to the "Red oak - mixed hardwood forest" type but with *Tsuga canadensis* (eastern hemlock) and/or *Pinus strobus* (eastern white pine) contributing more than 25% relative cover. Conifers may be scattered, locally abundant, may dominate the subcanopy, or may occur as a relict supracanopy (*Pinus strobus*), or in large former canopy gaps (*Pinus strobus*). *Quercus rubra* (northern red oak) is usually present, often dominant/codominant, most often with *Acer rubrum* (red maple), *Quercus velutina* (black oak), *Q. alba* (white oak), *Carya tomentosa* (mockernut hickory), *Betula lenta* (sweet birch), *Fraxinus americana* (white ash), *Fagus grandifolia* (American beech), and/or *Liriodendron tulipifera* (tuliptree). Shrubs include *Viburnum acerifolium* (maple-leaved viburnum), *Rhododendron periclymenoides* (pinxter-flower), *Amelanchier laevis* (smooth serviceberry), *A. arborea* (shadbush), *Carpinus caroliniana* (hornbeam), *Ostrya virginiana* (hop-hornbeam), *Hamamelis virginiana* (witch-hazel), and *Lindera benzoin* (spicebush). Herbaceous species include *Smilacina racemosa* (false Solomon's-seal), *Polygonatum biflorum* (Solomon's seal), *Gaultheria procumbens* (teaberry), *Maianthemum canadense* (Canada mayflower), and *Podophyllum peltatum* (may-apple).

Related types: The "Red oak - mixed hardwood forest" type has less than 25% combined relative cover by conifers. The type described here is generally dominated by a combination of various oaks—characteristically *Quercus rubra* (red oak), and *Tsuga canadensis* (eastern hemlock) and/or *Pinus strobus* (eastern white pine). In the "Hemlock (white pine) - northern hardwood forest," the same conifers usually share dominance with *Fagus grandifolia* (American beech), *Betula spp.* (birches), and *Acer saccharum* (sugar maple). The understory species associated with the "Hemlock (white pine) - northern hardwood forest" type are likewise more northern in affinity.

Range: Entire state except the Coastal Plain.

Selected references: Braun 1950, Eyre 1980.

[**Crosswalk:** Smith's "Dry - Mesic Acidic Central Forest" (in part), TNC's *Tsuga canadensis* - *Pinus strobus* Forest Alliance, SAF's White pine - northern red oak - red maple (20).]