

# Terrestrial & Palustrine Plant Communities Of Pennsylvania

By Jean Fike

## Rich hemlock - mesic hardwoods forest

These are species-rich, lower slope forests, reminiscent of the "Mixed mesophytic forest" type in the southwestern part of the state, but usually with a strong *Tsuga canadensis* (eastern hemlock) component. The hardwood species vary; typical representatives include *Liriodendron tulipifera* (tuliptree), *Fagus grandifolia* (American beech), *Quercus rubra* (northern red oak), *Acer rubrum* (red maple), *A. saccharum* (sugar maple), *Betula lenta* (sweet birch), *B. alleghaniensis* (yellow birch), *Fraxinus americana* (white ash), *Tilia americana* (basswood) and *Carya ovata* (shagbark hickory). Hemlock cover is often patchy. Under hardwood cover, the herbaceous diversity approaches that of the richer "Mixed mesophytic" type, while under dense hemlock cover, the herbaceous stratum reflects a more northern flora. *Magnolia tripetala*<sup>S</sup> (umbrella magnolia) is not uncommon. Other southern shrubs such as *Asimina triloba* (pawpaw) and *Staphylea trifolia* (bladdernut) may also occur, although *Rhododendron maximum* (rosebay), *Hamamelis virginiana* (witch-hazel), and *Lindera benzoin* (spicebush) are more abundant on most sites. Herbaceous species include *Adiantum pedatum* (maidenhair fern), *Erythronium americanum* (trout-lily), *Maianthemum canadense* (Canada mayflower), *Anemone quinquefolia* (wood anemone), *Dicentra canadensis* (squirrel-corn), *D. cucullaria* (dutchman's breeches), *Cimicifuga racemosa* (black snakeroot), *Geranium maculatum* (wood geranium), *Caulophyllum thalictroides* (blue cohosh), *Hepatica nobilis* (liverleaf), *Arisaema triphyllum* (jack-in-the-pulpit), *Allium tricoccum* (wild leek), *Sanguinaria canadensis* (bloodroot), *Corydalis flavula* (yellow fumewort), *Asplenium spp.* (spleenworts), *Botrychium virginianum* (rattlesnake fern), *Claytonia virginica* (spring-beauty), *Cardamine concatenata* (cut-leaved toothwort), *Mitella diphylla* (bishop's-cap), and *Asarum canadense* (wild ginger). In areas without a strong *Tsuga canadensis* (eastern hemlock) component, there may be complete annual litter turnover. This type may occur in a variety of lower slope/ravine situations.

**Related types:** This community type resembles a somewhat depauperate version of the "Mixed mesophytic forest" type, with the addition of *Tsuga canadensis* (eastern hemlock) usually with at least 25% relative cover. It is much richer in species composition than the most closely related mixed conifer/broadleaf forest type, the "Hemlock - tuliptree - birch forest." Species like *Magnolia tripetala* (umbrella magnolia), *Asimina triloba* (pawpaw), *Staphylea trifolia* (bladdernut), *Corydalis flavula* (yellow fumewort), *Sanguinaria canadensis* (bloodroot), and *Dicentra spp.* (dutchman's-breeches and squirrel-corn) are more typical of this richer, more southern type.

**Range:** Piedmont, Pittsburgh Plateau, southeastern portion of Ridge and Valley.

**Selected references:** Braun 1950, PNDI field surveys.

[**Crosswalk:** Smith's "Mesic Central Forest" (in part), TNC's *Tsuga canadensis* (mesic) Forest Alliance, SAF's Eastern hemlock (23) and Yellow poplar - white oak -northern red oak (59).]