

Green Floater (*Lasmigona subviridis*)

Freshwater Mussel Species of Concern

State Rank: S2 (imperiled), Global Rank: G3 (vulnerable)

Identification

The green floater (*Lasmigona subviridis*) is a small mussel, usually less than 55 mm in length. The shell is thin and the mussel has a subovate or trapezoidal shape. The color varies from a dull yellow to green with many dark green rays visible, especially in young individuals. This species may be confused with the creek heelsplitter (*Lasmigona compressa*) (NatureServe 2005; Strayer and Jirka 1997). The creek heelsplitter is larger, thicker shelled, and less ovate. Also, the creek heelsplitter has only been found in the Ohio River Drainage in Pennsylvania while the green floater is also present in the Susquehanna and Delaware River Drainages.



photo source: PNHP

Habitat

The green floater is often found in small creeks and large rivers and sometimes canals. This species is intolerant of strong currents and occurs in pools and other calm water areas (NatureServe 2005, North Carolina Mussel Atlas, Strayer and Jirka 1997). Preferred substrate is gravel and sand in water depths of one to four feet. This species is more likely to be found in hydrologically stable streams, not those prone to flooding and drying. Good water quality is also important for this mussel species (North Carolina Mussel Atlas).

Host Fish


Glochidial (larval) hosts for the green floater are not known (NatureServe 2005, Strayer and Jirka 1997).

Status

From New York south to Georgia and west to Tennessee the green floater is found. This species is not very common in Pennsylvania, but has been found in the Susquehanna, Delaware, and Ohio River Drainages (NatureServe 2005). The state status of the green floater is imperiled (S2), as it is not frequently encountered within its expected range (www.naturalheritage.state.pa.us/invertebrates.aspx). The small size of this species may make it difficult to locate live animals during surveys. Shells of dead green floaters tend to get buried in the surrounding habitat. More extensive surveys are necessary to determine the current status of this species in Pennsylvania and the United States.

The green floater was listed as threatened in an assessment of the conservation status of the freshwater mussels of the United States by the American Fisheries Society (Williams et al. 1993). The green floater has been historically widespread in the Susquehanna River drainage in New York; however, populations have declined since the early 1990s, probably due to pollution (Strayer and Jirka 1997). Decline in the abundance of this species in other places could be due to stream transport of their preferred habitat, as well as increases in pollutants. The introductions of zebra mussels and Asian clams have also negatively impacted abundance of this species in surveys. However, since this mussel species is hermaphroditic, small populations might survive slightly better than other mussel species in less than ideal conditions (NatureServe 2005).


Pennsylvania Distribution by County



Pennsylvania Natural Heritage Program data 2007

North American State/Province Conservation Status

Map by NatureServe (2007)



References

- NatureServe. 2007. NatureServe Explorer: An online encyclopedia of life [web application]. Version 6.2. NatureServe, Arlington, Virginia. Available <http://www.natureserve.org/explorer>. (Accessed: September 4, 2007).
- North Carolina Mussel Atlas, Species Information and Status. Website: www.ncwildlife.org/pg07_WildlifeSpeciesCon/pg7b1a1_9.htm
- Pennsylvania Natural Heritage Program. Biota of Concern In Pennsylvania (BOCIP) Lists. Website: www.naturalheritage.state.pa.us/invertebrates.aspx
- Strayer, D.L. and K.J. Jirka. 1997. The Pearly Mussels of New York State. The New York State Education Dept., Albany, NY 113pp and plates.
- Williams, J.D., M.L. Warren, K.S. Cummins, J.L. Harris, and R.J. Neves. 1993. Conservation Status of Freshwater Mussels. Fisheries 18(9): 6-22.

